

CHEMICALLY SYNTHESIZED haFGF GENE AND CORRESPONDING AMINO ACID SEQUENCE

FIG. 1

FGF fr HUMECGFB	TTC	GCC	CTG	ACC	AAG	AAT	CTG	CCT	CCA
haFGF 155	TTT	GCG	TTA	ACG	AAA	AAC	CTT	CCG	CCC
No amino acid	009	011	012	013	015	017	018	019	020
	Phe	Ala	Leu	Thr	Lys	Asn	Leu	Pro	Pro
FGF fr HUMECGFB	AAG	AAG	AAA	CTC	CTC	TGT	AGC	GGG	GGC
haFGF 155	AAA	AAA	AAG	CTT	CTT	TGC	AGT	GGA	GGA
No amino acid	024	025	027	028	029	031	032	034	035
	Lys	Lys	Lys	Leu	Leu	Cys	Ser	Gly	Gly
FGF fr HUMECGFB	AGG	ATC	CTT	CCG	GTG	ACA	AGG	GAC	AGG
haFGF 155	CGA	ATT	CTG	CCA	GTA	ACT	CGC	GAT	CGC
No amino acid	039	040	041	042	046	049	050	051	052
	Arg	Ile	Leu	Pro	Val	Thr	Arg	Asp	Arg
FGF fr HUMECGFB	AGC	CAG	AGT	GCG	GTG	GGG	GTG	ATA	AGT
haFGF 155	TCC	CAA	TCG	GCC	GTT	GGA	GTC	ATC	TCG
No amino acid	053	060	062	063	066	067	069	071	073
	Ser	Gln	Ser	Ala	Val	Gly	Val	Ile	Ser
FGF fr HUMECGFB	ACC	TTG	GAC	TTA	TAC	ACA	CCA	AAT	GAG
haFGF 155	ACG	CTT	GAT	CTG	TAT	ACG	CCT	AAC	GAA
No amino acid	074	080	085	088	089	093	094	095	096
	Thr	Leu	Asp	Leu	Tyr	Thr	Pro	Asn	Glu
FGF fr HUMECGFB	TGT	TTC	CTG	AGG	CTG	GAG	GAG	ACC	TAT
haFGF 155	TGC	TTT	CTA	AGA	CTA	GAA	GAA	ACG	TAC
No amino acid	098	100	101	103	104	105	106	111	112
	Cys	Phe	Leu	Arg	Leu	Glu	Glu	Thr	Tyr
FGF fr HUMECGFB	TCC	AAG	AAG	AAT	GTT	CTC	AAG	AAG	GGG
haFGF 155	TCG	AAA	AAA	AAC	GTA	CTT	AAA	AAA	GGT
No amino acid	114	115	116	121	124	126	127	128	130
	Ser	Lys	Lys	Asn	Val	Leu	Lys	Lys	Gly
FGF fr HUMECGFB	AGC	TGC	AAA	CGC	GGT	CCT	CAG	AAA	GCA
haFGF 155	TCC	TGT	AAG	CGT	GGA	CCA	CAA	AAG	GCT
No amino acid	131	132	133	134	135	136	142	143	144
	Ser	Cys	Lys	Arg	Gly	Pro	Gln	Lys	Ala
FGF fr HUMECGFB	TTT	CTC	CCC	CTG	GTC	TCT	TCT	GAT	
haFGF 155	TTC	CTG	CCA	CTC	GTG	AGC	TCC	GAC	
No amino acid	147	148	149	150	152	153	154	155	
	Phe	Leu	Pro	Leu	Val	Ser	Ser	Asp	

MODIFICATION IN MOLECULE haFGF 155 CODONS.

FGF fr HUMECGFB—THE SEQUENCE FROM GENBANK(at NCBI), haFGF 155
 —THE SEQUENCE SYNTHESIZED BY THE INVENTION AUTHORS.


FIG.2



FIG. 3

HPLC PURIFIED FGF155(8-23-00)

1 2 3



LANE1: 10 μ l OF CONDITIONED MEDIUM

2: 7 μ l OF HEPARIN-SEPHAROSE PURIFIED(0.45 μ g / μ l)

3: 14 μ l OUT OF 80 μ l OF HPLC-PURIFIED

FIG.4




FIG.5

NUCLEOTIDE AND AMINO ACID SEQUENCE FOR CHEMICALLY
SYNTHESIZED HUMAN A FGF (134 AMINO ACIDS)

FIG. 6



NUCLEOTIDE AND AMINO ACID SEQUENCE FOR CHEMICALLY
SYNTHESIZED HUMAN A FGF (140 AMINO ACIDS)

FIG. 8

9/12




FIG. 9

10/12

		BpiI	
2971	GGGCGCTGACTTCCGCGTTTCCAGACTTTACGAAACACGGAAACC CCCGCGACTGAAGGCGCAAAGGTCTGAAATGCTTTGTGCCTTTGG	GAAGACCATTTCATGTTGTTGCTCAGGTCGCAGACGTTTTGCAGCA CTTCTGGTAAGTACAACAACGAGTCCAGCGTCTGCAAAACGTCGT	
		PpuMI	
		EcoO109I	
		DraII	
		AvaII	
3061	GCACTCGCTTCACGTTTCGCTCGGTATCGGTGATTCTCTGCTA CGTCAGCGAAGTGCAAGCGAGCGCATAGCCACTAAGTAAGACGAT	ACCAGTAAGGCAACCCCGCCAGCCTAGCCGGGTCTCAACGACAG TGGTCATTCCGTTGGGGCGGTCTGGATCGGCCAGGAGTTGCTGTC	
		BsaBI	
		NsbI	
		BglII	
		VspI	
		BfmI	
		MbiI	
3151	GAGCAGCATCATGCGCACCCGTGGGGCCGCCAGATCTCGATCCCG CTCGTGCTAGTACGCGTGGGCACCCCGCGGTCTAGAGCTAGGGC	CGAAATTAATACGACTCACTATAGCGGAATTGTGAGCGGATAACA GCTTTAATTATGCTGAGTGATATCCCTTAACACTCGCCTATTGT	
		promotor	
		operator	
+2		MetAlaGluGlyGluIleThrThrPheThrAlaLeuThrGl	
		HpaI	
		HindII	
		HincII	
		XbaI	
		NdeI	
		Eco57I	
3241	ATTCCCTCTAGAAATAATTTTGTTTAACTTTAAGAAGGAGATAT TAAGGGGAGATCTTTATTAACAAATTGAAATCTCTCTCTATA	ACATATGGCTGAAGGGGAAATCACCACCTTTACAGCGTTAACGGA TGTATACCGACTTCCCTTTAGTGGTGAATGTGCGAATTGCT	
+2	uLysPheAsnLeuProProGlyAsnTyrLysLysProLysLeuLeu	TyrCysSerAsnGlyGlyHisPheLeuArgIleLeuProAspGl	
		SmaI	
		PstI	
		AvaI	
		HindIII	
		BfmI	
		EcoRI	
3331	GAAATTTAACCTTCCGCCCGGGAATTACAAAAACCCAGCTTCT CTTTAAATTGGAAGCGGGCCCTTAATGTTTTTGGGTTCGAAGA	TTACTGCAGTAACGGAGGACACTTCCTGCGAATTCTGCCAGATGG AATGACGTCATTGCCTCCTGTGAAGGACGCTTAAGACGGTCTACC	
+2	yThrValAspGlyThrArgAspArgSerAspGlnHisIleGlnLeu	GlnLeuSerAlaGluSerValGlyGluValTyrIleLysSerTh	
		SalI	
		HindII	
		HincII	
		PvuI	
		XmaIII	
		NruI	
		PvuII	
		CfrI	
3421	CACAGTAGATGGGACTCGCGATCGCTCCGACCAGCACATTACGCT GTGTCATCTACCCTGAGCGCTAGCGAGGCTGGTGTGAAGTCGA	GCAACTCTCGGCCGAAAGCGTTGGAGAGGTCTATATCAAGTTCGAC CGTTGAGAGCCGGCTTCGCAACCTCTCCAGATATAGTTCAGCTG	
+2	rGluThrGlyGlnTyrLeuAlaMetAspThrAspGlyLeuLeuTyr	GlySerGlnThrProAsnGluGluCysLeuPheLeuGluArgLe	
		MlsI	
		RsaI	
		StyI	
		CfrI	
		Csp6I	
		NcoI	
		AcyI	
		MvaI269I	
		XbaI	
3511	GGAGACTGGCCAGTACCTTGCCATGGACACCGATGGGCTTCTGTA CCTCTGACCGGTCATGGAACGGTACCTGTGGTACCCGAAGACAT	TGGCTCACAGACGCCTAACGAAGAATGCTTGTCTTAGAAAGACT ACCGAGTGTCTGCGGATTGCTTCTTACGAACAAAGATCTTTCTGA	
		BsmAI	

NUCLEOTIDE AND AMINO ACID SEQUENCE FOR CHEMICALLY
SYNTHESIZED INTERFERON α -2B

FIG. 10



1-MOLECULAR WEIGHT KIT(94 000,67 000,43 000,30 000
20 100, 14 400)

2-haFGF 134(40 μ I OF THE CULTURAL MEDIUM)

3-haFGF 140(40 μ l OF THE CULTURAL MEDIUM)

4-IFN α 2B(40 μ l OF THE CULTURAL MEDIUM)


5-haFGF 155(40 μ l OF THE CULTURAL MEDIUM)

6-HGH(40 μ l OF THE CULTURAL MEDIUM)

7-MAP(40 μ l OF THE CULTURAL MEDIUM)

8- β -GALACTOSIDASE OF E. COLI(40 μ l OF THE CULTURAL MEDIUM)

FIG. 11



ELECTROPHOREGRAMM OF THE PURIFIED PRODUCTS:

1-MOLECULAR WEIGHT KIT (94 000, 67 000, 43 000, 30 000, 20 100, 14 400)

2-haFGF 134

3-haFGF 140

4-haFGF 146

5-IFN α 2b

6-haFGF 155

7-MAP

8-MOLECULAR WEIGHT KIT

FIG. 12